

Gravitational Scattering in the ADD-model Revisited

Malin Sjö Dahl

Dept. of Theoretical Physics, Sölvegatan 14A, S-223 62 Lund, Sweden

E-mail: Malin.Sjodahl@thep.lu.se

ABSTRACT: The effects of the condition that the standard model particles live on a thin brane in the ADD-model is investigated. This is shown to result in a cut-off independent Kaluza–Klein propagator for large distance scattering and an odd number of extra dimensions. The matrix element corresponding to this propagator can also be Fourier transformed to position space, giving back the extra-dimensional version of Newton’s law. For an even number of extra dimensions the corresponding propagator is found by requiring that Newton’s law should be recovered.

KEYWORDS: ADD, Extra dimensions, Gravitational scattering.

Contents

1. Introduction	1
2. Fourier transformation to position space in the extra dimensions	3
3. Fourier transformation to position space in our ordinary dimensions	4
4. Conclusion and outlook	5

1. Introduction

The ADD-model [1–3] aims at explaining the hierarchy between the weak scale and the Planck scale. This is done by introducing extra, compactified, dimensions in which only gravity is allowed to propagate. At distances small compared to the compactification radius, but large compared to the Planck length and the thickness of the brane, the gravitational force will be enhanced and behave essentially as in a $4 + n$ dimensional world, where n is the number of extra dimensions. If the extra dimensions are large enough, the enhanced gravitational force opens up for the possibility of gravitational scattering and black hole production at present, or soon upcoming, collider experiments.

To quantify the amount of gravitational interaction, the theory was put on a perturbative field-theoretical basis in [4, 5]. The perturbations of the extra dimensional part of the metric enter as massive Kaluza–Klein (KK) modes in the Lagrangian. When these modes are internal states they have to be summed over which lead the authors of [4, 5] to the following divergent propagator integral

$$\sum_{\vec{l}} \frac{1}{-m_{\vec{l}}^2 + k^2} \sim R^n \int \frac{m^{n-1}}{-m^2 + k^2} dm. \quad (1.1)$$

Here $\vec{l}$ enumerates the allowed momenta, $m_{\vec{l}}$, in the extra dimensions, m is the absolute value of $m_{\vec{l}}$, R is the compactification radius and k^2 is the momentum squared of the $3 + 1$ -dimensional part of the propagator. (This object, without Lorentz structure, will somewhat sloppily be referred to as a propagator.) For the above approximation to be valid, the compactification radius R of the extra dimensions clearly have to be large compared to other relevant length scales. As it stands, the integral eq. (1.1) is explicitly divergent for $n \geq 2$. However, when arriving at eq. (1.1), the physical condition that the standard model fields are confined to the brane was not taken into account.

Instead the problem of the divergent integral was approached in [4, 5] by introducing a cut-off M_s , argued to be of the same order of magnitude as the fundamental Planck scale, M_p . (A physical motivation for a cut-off was later considered in [6, 7] by the introduction

of a brane tension, and various cut-off versions have been discussed in [8]). For $n > 2$, and exchanged momentum small compared to M_s , the Kaluza–Klein summation of t-channel (or s-channel) amplitudes then gave a propagator behaving as

$$\frac{1}{n-2} R^n M_s^{n-2} \sim \frac{1}{G_{N(4)}} \frac{1}{n-2} \frac{M_s^{n-2}}{M_p^{n+2}} \quad (1.2)$$

where $G_{N(4)}$ is the ordinary 3+1-dimensional Newton’s constant related to R and M_p via $G_{N(4)}^{-1} \approx R^n M_p^{n+2}$. (In [4] M_p and M_s are taken to be the same in the calculation of the propagator.) In the Born approximation the cross section would then be given by [9]

$$\frac{d\sigma}{dz} \sim \frac{s^3}{(n-2)^2} \left(\frac{M_s^{n-2}}{M_p^{n+2}} \right)^2 F(\text{spin}, z), \quad (1.3)$$

where z is the cosine of the scattering angle in the center of mass system, s the squared sum of the incoming particles momenta and F a function taking spin dependence into account.

A different approach to calculating the influence of the Kaluza–Klein modes, was presented in [2] when deriving Newton’s law. In this case the summation of Kaluza–Klein modes was performed in the classical limit, *after* Fourier transforming our normal momentum space to coordinate space [2], giving rise to the expected $3 + n$ dimensional version of Newton’s law,

$$\frac{V(r)}{m_1 m_2} \sim \int_{-\infty}^{\infty} d\bar{m} \int_{-\infty}^{\infty} d\bar{k} \frac{1}{\bar{m}^2 + \bar{k}^2} e^{i\bar{k} \cdot \bar{x}} \sim \int_0^{\infty} dm m^{n-1} \frac{e^{-mr}}{r} \sim \frac{1}{r^{n+1}} \quad (1.4)$$

where $r = |\bar{x}|$. Eq. (1.2) and eq. (1.4) may seem to contradict each other. Especially, eq. (1.2) gives the same form of the gravitational scattering regardless of the number of extra dimensions, namely a δ -function at $r = 0$. Equation (1.4), on the other hand, gives different scattering behavior for different number of extra dimensions. We will see in the next section that k -dependent correction terms to eq. (1.2) are important for the classical limit as the $1/r^{n+1}$ -potential can not be recovered by Fourier transforming the non-relativistic matrix element corresponding to eq. (1.2) to ordinary position space.

As we like to implement the condition that standard model fields live on a brane via a Fourier transform, we should really add a factor $i\bar{y} \cdot \bar{m}$, where $\bar{y}$ is the coordinate in the extra dimensions, in the exponent and then integrate over a narrow distribution in $\bar{y}$. However, as the m integral is explicitly convergent when evaluated after the k integral, the addition of $i\bar{y} \cdot \bar{m}$ in the exponent would not change the result in the classical limit when the distance is much larger than the brane thickness. In essence, what will be done in this paper is nothing but calculating eq. (1.4) in the reversed order, starting with the m -integral.

The Fourier transform to position space w.r.t. the extra dimensions is performed (for an odd number of extra dimensions) in section 2. In section 3 we show that the large distance limit of the resulting propagator can be further Fourier transformed to position space w.r.t. our normal dimensions giving back the extra-dimensional version of Newton’s law. We also find the corresponding KK-”summed” propagator for an even number of extra

dimensions by requiring that Newton's law should be recovered. Finally we summarize and conclude in section 4.

2. Fourier transformation to position space in the extra dimensions

As the standard model fields are assumed to live on the brane, any measurement of gravitational scattering will be in position space w.r.t. the extra dimensional coordinates and in momentum space w.r.t. our ordinary coordinates. We therefore search for the corresponding propagator. To find it we introduce a coordinate $\bar{y}$, with absolute value y , in the extra dimensions. Later we will be interested in a narrow distribution around $\bar{y} = \bar{0}$ corresponding to a small extension of the standard model fields into the bulk. Searching for a propagator which is in position space w.r.t. the extra dimensions also corresponds to imposing the condition of locality. As the standard model fields live on a brane they will only be sensitive to KK modes which overlap with the brane.

Fourier transforming eq. (1.1) to position space with respect to the extra dimensions now gives for $n \geq 3$

$$\begin{aligned} D(k, y) &= \sum_{\vec{l}} \frac{1}{-m_{\vec{l}}^2 + k^2} e^{i2\pi\vec{l}\cdot\bar{y}/R} \\ &\approx \left(\frac{R}{2\pi}\right)^n S_{n-1} \int_0^\pi \sin(\theta)^{n-2} d\theta \int_0^\infty dm \frac{m^{n-1}}{-m^2 + k^2} e^{imy \cos(\theta)} \\ &= \left(\frac{R}{2\pi}\right)^n S_{n-1} \int_{-1}^1 d\cos(\theta) (1 - \cos(\theta)^2)^{(n-3)/2} \int_0^\infty dm \frac{m^{n-1} e^{imy \cos(\theta)}}{-m^2 + k^2} \quad (2.1) \end{aligned}$$

where S_{n-1} is the surface of a unit sphere in $n - 1$ dimensions (from integration over the angles on which the integrand does not depend) and the factor $R^n/(2\pi)^n$ comes from the density of Kaluza–Klein modes. For an odd number of extra dimensions this can, with $x = \cos(\theta)$, be rewritten as

$$\begin{aligned} D(k, y) &= \left(\frac{R}{2\pi}\right)^n S_{n-1} \int_{-1}^1 dx (1 - x^2)^{(n-3)/2} \int_0^\infty dm \\ &\quad \left(- \sum_{j=0}^{(n-3)/2} m^{n-3-2j} k^{2j} + \frac{k^{n-1}}{-m^2 + k^2} \right) e^{imyx}. \quad (2.2) \end{aligned}$$

From this form we see that the terms in the sum are either k -independent, and therefore correspond to δ -functions at $r = 0$ in ordinary position space, or are even powers of k and correspond to derivatives of the δ -function. (This is easily seen by Fourier transforming from k -space to r -space component by component.) These terms are therefore localized to distance zero in ordinary space and can not be important in the classical region, where wave packages are well separated and do not overlap.

If we imagine the brane to have a finite narrow distribution (such as a Gaussian) around $\bar{y} = \bar{0}$ in the extra dimensions, the $1/r^{n+1}$ -law will be modified at distances similar to the brane thickness. We also note that the terms in the sum give (possibly) large but finite

contributions. These terms, which depend on the extension of the brane, are important when the wave functions overlap and may then give an interaction similar to that from eq. (1.3) (until the unitarity condition sets in) but they will not be further investigated here. Instead we concentrate on the term relevant for large distance and large energy scattering (the classical limit) coming from the last term in the integral. The contribution to the integral from this term, called $\hat{D}(k)$, is easily evaluated in the limit of small y (corresponding to a narrow distribution) and is given by

$$\hat{D}(k) \approx \left(\frac{R}{2\pi}\right)^n \frac{\pi S_n}{2} \left(\sqrt{-k^2}\right)^{n-2} i^{n+1}. \quad (2.3)$$

It is easy to show that this result holds also for one extra dimension, and therefore for any odd number of extra dimensions.

The expression eq. (2.3), which is in momentum space with respect to our normal dimensions, and in position space with respect to the extra dimensions, has the following properties:

- i It gives back Newton's law, eq. (1.4). This will be demonstrated in section 3.
- ii It depends on the number of extra dimensions in a non-trivial way, such that, as the gravitational force increases faster with smaller distance in position space for many extra dimensions, this is reflected in a faster increase with larger k in momentum space.
- iii It does not depend on an arbitrary cut-off as long as the cut-off ($\approx 1/(\text{brane thickness})$) is much larger than k . This implies that it is not dominated by metric perturbations of the scale $1/M_p$ for scatterings corresponding to much larger distances. This is the case for eq. (1.1) integrated to $M_s \approx M_p$.
- iiii It is the part of the propagator which is argued to contribute to the all order exponentiated eikonalized amplitude in [10] (apart from what appears to be a sign misprint).

3. Fourier transformation to position space in our ordinary dimensions

To obtain the 3+1+n dimensional version of Newton's law we take the classical limit such that the energy is given by the mass, multiply with the coupling constant $4\pi G_{N(4)}$ and Fourier transform eq. (2.3) to position space. Using $\kappa = |\vec{k}|$ we have

$$\begin{aligned} \frac{V(r)}{m_1 m_2} &= 4\pi G_{N(4)} \left(\frac{R}{2\pi}\right)^n \frac{\pi S_n}{2} i^{n+1} \frac{S_3}{2} \frac{1}{(2\pi)^3} \int_{-1}^1 d\cos(\theta) \int_0^\infty d\kappa \kappa^2 \kappa^{n-2} e^{i\kappa r \cos(\theta)} \\ &= 4\pi G_{N(4)} \left(\frac{R}{2\pi}\right)^n \frac{\pi S_n S_3}{4(2\pi)^3} i^{n+1} \frac{1}{ir} \int_0^\infty d\kappa \kappa^{n-1} (e^{i\kappa r} - e^{-i\kappa r}). \end{aligned} \quad (3.1)$$

This can be evaluated by introduction of a small convergence factor

$$\frac{V(r)}{m_1 m_2} = 4\pi G_{N(4)} \left(\frac{R}{2\pi}\right)^n \frac{\pi S_n S_3}{4(2\pi)^3} i^{n+1} \frac{1}{ir} \lim_{\epsilon \rightarrow 0} \left(\frac{d}{dr}\right)^{n-1} \int_0^\infty d\kappa \left[\frac{e^{i\kappa r - \epsilon \kappa}}{i^{n-1}} - \frac{e^{-i\kappa r - \epsilon \kappa}}{(-i)^{n-1}} \right]$$

$$\begin{aligned}
&= G_{N(4)} \left(\frac{R}{2\pi} \right)^n \frac{\pi S_n (4\pi)^2}{4(2\pi)^3} i^{n+1} \frac{1}{ir} \frac{1}{i^{n-1}} \lim_{\epsilon \rightarrow 0} \left(\frac{d}{dr} \right)^{n-1} (-1) \left[\frac{-i}{r+i\epsilon} - \frac{i(-1)^{n-1}}{r-i\epsilon} \right] \\
&= -G_{N(4)} \left(\frac{R}{2\pi} \right)^n S_n \frac{\Gamma(n)}{r^{n+1}}, \tag{3.2}
\end{aligned}$$

where the last step only is valid for an odd number of extra dimensions. This is the same result as in [2] (apart from a minus sign which is neglected in [2]), i.e. gravitational scattering enhanced by the large density of Kaluza–Klein modes, corresponding to a large coupling constant.

The strategy so far has been to start from the propagator and argue that we can get back Newton’s law. Clearly this argument could be turned upside down. Using the result in eq. (1.4) [2], we could alternatively search for the propagator giving the expected potential when Fourier transformed to position space. Again this would give us a propagator of the form k^{n-2} for an odd number of extra dimensions. For an even number of extra dimensions we settle with this argument, showing that the propagator

$$\hat{D}(k) \approx \left(\frac{R}{2\pi} \right)^n \frac{S_n}{2} (-1)^{\frac{n-2}{2}} \left(\sqrt{-k^2} \right)^{n-2} \ln(-k^2) \tag{3.3}$$

gives the desired form. The logarithm of a dimension-full quantity may seem disturbing. Replacing it by $\ln(-k^2/k_0^2)$ for some k_0 we note that the $k^{n-2} \ln(k_0^2)$ -term would only contribute at $r = 0$ when Fourier transformed to position space. This means that, just as in the case of an odd number of extra dimensions, we have a local contribution which depends on the brane. We also note that, just as for odd n , requiring the standard model particles to live on a thin brane by introducing a (narrow) distribution in y -space we would have a (wide) distribution in m -space giving a (large) finite value for the propagator.

Fourier transforming the non-relativistic version of eq. (3.3) to position space we find the potential

$$\begin{aligned}
\frac{V(r)}{m_1 m_2} &= 4\pi G_{N(4)} \left(\frac{R}{2\pi} \right)^n \frac{S_n}{2} (-1)^{\frac{n-2}{2}} \frac{S_3}{2} \frac{1}{(2\pi)^3} \int_{-1}^1 d\cos(\theta) \int_0^\infty d\kappa \kappa^2 \kappa^{n-2} \ln(\kappa^2) e^{i\kappa r \cos(\theta)} \\
&= G_{N(4)} \left(\frac{R}{2\pi} \right)^n \frac{S_n (4\pi)^2}{4(2\pi)^3} (-1)^{\frac{n-2}{2}} \frac{1}{ir} \int_0^\infty d\kappa \kappa^{n-1} \ln(\kappa^2) 2i \sin(\kappa r). \tag{3.4}
\end{aligned}$$

For n even we get after evaluating the integral [11]

$$\frac{V(r)}{m_1 m_2} = -G_{N(4)} \left(\frac{R}{2\pi} \right)^n S_n \frac{\Gamma(n)}{r^{n+1}}. \tag{3.5}$$

This is the multidimensional version of Newton’s law. We also note that eq. (3.3) is the part of the propagator argued to contribute to the all order eikonalized amplitude in [10].

4. Conclusion and outlook

We have shown that requiring the standard model particles to live on a finite brane, leads to a convergent result for the KK propagator. The part of the propagator which is relevant

for large energy and large distance scattering can also (at least for an odd number of extra dimensions) be Fourier transformed to position space w.r.t. our ordinary coordinates, giving back Newton’s law. For an even number of extra dimensions we have found an expression for the KK-”summed” propagator by requiring that we should get back Newton’s law in the classical limit, when all coordinates are Fourier transformed to position space.

It should be pointed out that, although the tree-level amplitude is calculated to a finite value, we can not use eq. (2.3) and eq. (3.3) for calculating large energy cross sections with the Born approximation. Instead an all order summation still has to be performed [10]. Furthermore, the cross section obtained in [10] is supported by the results here, as the finite results for the propagator here agrees with the parts argued to contribute to the eikonal cross section in [10].

Acknowledgments

I am thankful to Leif Lönnblad, Johan Grönqvist, Gösta Gustafson, Pavel Kurasov and Johan Bijnens for useful discussions and comments on the manuscript and to Gian Giudice for making me aware of some of the cited articles.

References

- [1] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali *Phys. Lett.* **B429** (1998) 263–272, [hep-ph/9803315](#).
- [2] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali *Phys. Rev.* **D59** (1999) 086004, [hep-ph/9807344](#).
- [3] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali *Phys. Lett.* **B436** (1998) 257–263, [hep-ph/9804398](#).
- [4] T. Han, J. D. Lykken, and R.-J. Zhang *Phys. Rev.* **D59** (1999) 105006, [hep-ph/9811350](#).
- [5] G. F. Giudice, R. Rattazzi, and J. D. Wells *Nucl. Phys.* **B544** (1999) 3–38, [hep-ph/9811291](#).
- [6] M. Bando, T. Kugo, T. Noguchi, and K. Yoshioka *Phys. Rev. Lett.* **83** (1999) 3601–3604, [hep-ph/9906549](#).
- [7] T. Kugo and K. Yoshioka *Nucl. Phys.* **B594** (2001) 301–328, [hep-ph/9912496](#).
- [8] G. F. Giudice and A. Strumia *Nucl. Phys.* **B663** (2003) 377–393, [hep-ph/0301232](#).
- [9] D. Atwood, S. Bar-Shalom, and A. Soni *Phys. Rev.* **D62** (2000) 056008, [hep-ph/9911231](#).
- [10] G. F. Giudice, R. Rattazzi, and J. D. Wells *Nucl. Phys.* **B630** (2002) 293–325, [hep-ph/0112161](#).
- [11] I. M. Gradshteyn and I. Ryshik, *Table of Integrals Series, and Products*. Academic Press, USA, 1994.